

- their relationship to human isolates. *J Antimicrob Chem.* 2006; 57(3): 461-5.
2. Wijaya L, Hsu LY. Community-associated methicillin-resistant *Staphylococcus aureus* skin and soft tissue infections. *Pro Sing Healthcare.* 2010; 19(3): 212-9.
 3. Grisold AJ, Leitner E, Muhlbauer G, et al. Detection of methicillin-resistant *Staphylococcus aureus* and simultaneous confirmation by automated nucleic acid extraction and real-time PCR. *J Clin Microbiol.* 2010; 40(7): 2392-7.
 4. Tiemersma EW, Bronzwaer SL, Lyytikainen O. Methicillin-resistant *Staphylococcus aureus* in Europe 1999-2002. *Emerging Inf Dis.* 2010; 10(5): 1627-34.
 5. Chambers HF. Detection of methicillin-resistant staphylococci. *Inf Dis Clin Nor Am.* 1993; 7(1): 425-33.
 6. Hartman BJ, Tomasz A. Low-affinity penicillin-binding protein associated with β -lactam resistance in *Staphylococcus aureus*. *J Bactiol.* 1984; 158(8): 513-6.
 7. Waxman DJ, Strominger JL. Penicillin-binding proteins and the mechanism of action of beta-lactam antibiotics. *Annu Rev Biochem.* 1983; 52(4): 825-69.
 8. Kluytmans J, van Belkum A, Verbrugh H. Nasal carriage of *Staphylococcus aureus*: epidemiology underlying mechanisms and associated risks. *Clin Microbiol Rev.* 1997; 10(3): 505-20.
 9. Miller LG, Diep BA. Colonization Fomites and Virulence: Rethinking the Pathogenesis of Community-Associated Methicillin-Resistant *Staphylococcus aureus* Infection. *Clin Inf Dis.* 2008; 46(1): 752-60.
 10. Maltezou, HC Giamarellou, H. Community-acquired methicillin-resistant *Staphylococcus aureus* infections. *Int J Antimicrob Agents.* 2006; 11(2): 112-20.
 11. Naimi TS, LeDell KH, Como-Sabetti K, et al. Comparison of community and healthcare-associated methicillin-resistant *Staphylococcus aureus* infection. *J Am Med Assoc.* 2003; 290(6): 2976-84.
 12. Herold BC, Immergluck LC, Maranan MC, et al. Community acquired methicillin-resistant *Staphylococcus aureus* in children with no identified predisposing risk. *J Am Med Assoc.* 2008; 279(10): 593-8.
 13. Ho PL, Chuang SK, Choi YF, et al. Community-associated methicillin-resistant and methicillin-sensitive *Staphylococcus aureus*: skin and soft tissue infections in Hong Kong. *Diagn Microbiol Infect Dis.* 2008; 61(7): 245-50.
 14. Chen CJ, Hsueh PR, Su LH, et al. Change in the molecular epidemiology of methicillin-resistant *Staphylococcus aureus* bloodstream infections in Taiwan. *Diagn Microbiol Infect Dis.* 2004; 65(4): 199-201.
 15. Gonzalez BE, Rueda AM, Shelburne SA, et al. Community-associated strains of methicillin-resistant *Staphylococcus aureus* as the cause of healthcare-associated infection. *Infect Control Hosp Epidemiol.* 2006; 27(1): 1051-6.
 16. Maree CL, Daum RS, Boyle-Vavra S, et al. Community-associated methicillin-resistant *Staphylococcus aureus* isolates causing healthcare associated infections. *Emerging Infect Dis.* 2007; 13(3): 236-42.
 17. Bartels MD, Kristoffersen K, Boye K, et al. Rise and subsequent decline of community-associated methicillin-resistant *Staphylococcus aureus* ST30-IVc in Copenhagen Denmark through an effective search and destroy policy. *Clin Microbiol Infect.* 2007; 16(10): 78-83.
 18. Miller LG, Perdreau-Remington F, Rieg G, et al. Necrotizing fasciitis caused by community-associated methicillin-resistant *Staphylococcus aureus* in Los Angeles. *New Eng J Med.* 2005; 352(6): 1445-53.
 19. Lee YT, Lin JC, Wang NC, et al. Necrotizing fasciitis in a medical center in northern Taiwan: emergence of methicillin-resistant *Staphylococcus aureus* in the community. *J Microbiol Immunol Infect.* 2007; 40(2): 335-41.
 20. Martinez-Aguilar G, Avalos-Mishaan A, Hulten K, et al. Community-acquired methicillin-resistant and methicillin-susceptible *Staphylococcus aureus* musculoskeletal infections in children. *Pediatr Infect Dis J.* 2004; 23(2): 701-6.
 21. Pannaraj PS, Hulten KG, Gonzalez BE, et al. Infective pyomyositis and myositis in children in the era of community-acquired methicillin-resistant *Staphylococcus aureus* infection. *Clin Inf Dis.* 2006; 43(4): 953-60.
 22. Francis JS, Doherty MC, Lopatin U, et al. Severe community-onset pneumonia in healthy adults caused by methicillin-resistant *Staphylococcus aureus* carrying the Panton-Valentine leukocidin genes. *Clin Inf Dis.* 2005; 40(9): 100-7.
 23. Centers for Disease Control and Prevention from the Centers for Disease Control and Prevention for pediatric deaths from community-acquired methicillin-resistant *Staphylococcus aureus* - Minnesota and North Dakota 1997-1999. *J Am Med Assoc.* 2004; 282: 1123-5.
 24. Buck JM, Como-Sabetti K, Harriman KH, et al. Community-associated methicillin-resistant *Staphylococcus aureus* Minnesota 2000-2003. *Emerging Inf Dis.* 2005; 11(5): 1532-8.
 25. Fridkin SK, Hageman JC, Morrison M, et al. Methicillin-resistant *Staphylococcus aureus* disease in three communities. *New Engl J Med.* 2005; 352(11): 1436-44.
 26. Frazee BW, Lynn J, Charlebois ED, et al. High prevalence of methicillin-resistant *Staphylococcus aureus* in emergency department skin and soft tissue infections. *Ann Emerg Med.* 2005; 45(6): 311-20.
 27. Kateete DP, Kimani CN, Katabazo FA, et al. Identification of *Staphylococcus aureus*: DNase and Mannitol salt agar improve the efficiency of the tube coagulase test. *Ann Clin Microbiol Antimicro.* 2010; 9(23): 612-20.