

28. Rosana BR, Natalia LP, Iorio Malvar KL, et al. Coagulase-negative Staphylococci: comparison of phenotypic and genotypic oxacillin susceptibility tests and evaluation of the agar screening test by using different concentrations of oxacillin. J Clin Microbiol. 2003; 41(8): 3609-14.
29. Clinical and Laboratory Standards Institute. Performance standards for antimicrobial disk susceptibility tests Approved standard M02-A10 WaynePA: Clinical and Laboratory Standards Institute, 2009.
30. Lecercq R, Bauduret F, Soussy C. Selection of constitutive mutants of gram-positive cocci inducible resistant to macrolides lincosamides and streptogramins (MLS): comparison of the selective effects of the MLS. Pathol Biol. 1989; 37(10): 568-72.
31. Trieu-Cuot P, Poyart-Salmeron C, Carlier C, et al. Nucleotide sequences of the erythromycin resistant gene of the conjugative transposon Tn1545. Nucleic Acids Res. 1990; 18(3): 3660-70.
32. Ross JI, Eady EA, Cove JH, et al. Inducible erythromycin resistance in staphylococci is encoded by a member of ATP-binding transport super-gene family. Mol Microbiol. 1990; 4(7): 1207-14.
33. Brisson-Noël A, Delrieu P, Samain D, et al. Inactivation of lincosamides antibiotics in Staphylococcus identification of lincosamide o-nucleotidyltransferases and comparison of the corresponding resistance gene. J Biol Chem. 1988; 263(4): 15880-7.
34. Ardic N, Ozyurt M, Sareyyupoglu B, et al. Investigation of erythromycin and tetracycline resistant genes in methicillin-resistant staphylococci. Int J Antimicrob Agent. 2005; 26 (12): 213-8.
35. Rushdy AA, Salama MS, Othman AS. Detection of methicillin/oxacillin resistant *Staphylococcus aureus* isolated from some clinical hospital in Cairo using MecA/Nuc gene and antibiotic susceptibility profile. Int J Agricul Biol. 2007; 9(6): 211-7.
36. Jaffe RI, Lane JD, Albury SV, et al. Rapid extraction from and direct identification in clinical samples of methicillin-resistant staphylococci using PCR. J Clin Microbiol. 2000; 38(7): 3407-12.
37. Huletsky A, Giroux R, Rossbach V, et al. New Real-Time PCR Assay for Rapid Detection of Methicillin-Resistant *Staphylococcus aureus* Directly from Specimens Containing a Mixture of Staphylococci. J Clin Microbiol. 2004; 42(5): 1874-84.
38. Thakker-varia S, Jenssen WD, Moon-McDermott L, et al. Molecular epidemiology of macrolide-lincosamide-streptogramin B resistance in *Staphylococcus aureus* and coagulase-negative staphylococci. Antimicrob Agents Chemother. 1987; 31(10): 735-43.
39. Eady EA, Ross KI, Tipper JL, et al. Distribution of genes encoding erythromycin ribosomal methylases and an erythromycin efflux pump in epidemiologically distinct groups of staphylococci. Antimicrob Agents Chemother. 1993; 31(2): 211-7.
40. Leclercq R, Brisson-Noël A, Duval J, et al. Phenotypic expression and gene heterogeneity of lincosamide inactivation in *Staphylococcus* spp. Antimicrob Agents Chemother. 1987; 31(7): 1887-91.
41. Francois P, Pittet D, Bento M, et al. Rapid detection for methicillin-resistant *Staphylococcus aureus* directly from sterile or nonsteril clinical samples by a new molecular assay. J Clin Microbiol. 2003; 41(1): 254-60.
42. Horman MA, Atoom AM, Abuharfeil NM, et al. Identification of methicillin-resistant *Staphylococcus aureus* and methicillin-resistant coagulase-negative staphylococci in clinical setting. Am J Inf Dis. 2008; 4(2): 156-61.
43. National Standard Method Coagulase SOP UK. (<http://www.hpa-standardmethods.org.uk/>).
44. Cappuccino JG, Sherman N. Microbiology: A Laboratory Manual 8<sup>th</sup> ed. PEARSON: Benjamin Cummings, 2008.

---

Correspondence to Dr. Ahmed S Abdulmir

E-mail: Ahmsah73@yahoo.com

Received 11<sup>th</sup> Dec. 2011: Accepted 9<sup>th</sup> Feb. 2012