

2. Sterns EE. Age-related breast diagnosis. *Can J Surg.* 1992; 35: 41-5.
3. Ljung BM, Drejet A, Chiampi N, et al. Diagnostic accuracy of fine-needle aspiration biopsy is determined by physician training in sampling technique. *Cancer.* 2001; 93:263-8.
4. Al-Hasnawi S, Alkhuza'ie A, Al-Mosawi AJ, et al. The histopathologic pattern of breast cancer in Iraq. *The N Iraqi J Med.* 2009; 5(2): S77-9.
5. Zayicek J. Aspiration biopsy in diagnosis of palpable mammary lesions. In: Wied GL, Koss LG, Reagan JW (eds). *Compendium of diagnostic cytology.* Vol. IV, University of Chicago. 1976; p. 555-75.
6. Donegan WL. Evaluation of a palpable breast mass. *N Engl J Med.* 1992; 327: 937-42.
7. Sickle EA, Parker SH. Appropriate role of core breast biopsy in the management of probably benign lesions. *Radiology.* 1993 Aug; 188(2): 315.
8. Dawson AE, Mulford DK, Taylor AS, et al. Breast carcinoma in women age 35 years and younger: mammography and diagnosis by fine-needle aspiration cytology. *Cancer.* 1998; 84: 163-8.
9. American College of Radiology. *Breast imaging reporting and data system: breast imaging atlas*, 4th ed. Reston, VA: American College of Radiology, 2003.
10. Oswald G, Helbich TH, Fuchsjaeger MH, et al. Follow-up of Palpable Circumscribed Non-calcified Solid Breast Masses at Mammography and US: Can Biopsy Be Averted? *Radiology.* 2004; 233(3): 850-6.
11. Oswald G, Helbich TH, Hopf G, et al. Probably benign breast masses at US: is follow-up an acceptable alternative to biopsy? *Radiology.* 2007; 244: 87-93.
12. Tabar L, Duffy SW, Vitak B, et al. The natural history of breast carcinoma: what have we learned from screening? *Cancer.* 1999; 86: 449-62.
13. Kocjan G. Needle aspiration cytology of the breast: current perspective on the role in diagnosis and management. *Acta Med Croatica.* 2008 Oct; 62(4): 391-401.
14. Kerlikowske K, Smith-Bindman R, Ljung BM, et al. Evaluation of Abnormal Mammography Results and Palpable Breast Abnormalities. *Ann Intern Med.* 2003 Aug 19; 139(4): 274-84.
15. Rosenthal DL. Breast lesions diagnosed by FNAC. *Path Res Pract Dec.* 1986; 381(6): 645-56.
16. Lieske B, Ravichandran D, Wright D. Role of fine-needle aspiration cytology and core biopsy in the preoperative diagnosis of screen-detected breast carcinoma. *Br J Cancer.* 2006 July 3; 95(1): 62-6.
17. Nooraddin I. Fine needle aspiration cytology (FNAC) versus open biopsy in breast masses in Erbil. *The N Iraqi J Med.* 2008; 4(2): 15-21.
18. Parkin DM, Bray F, Ferlay J, et al. Estimating the world cancer burden: Globocon 2000. *Int J Cancer.* 2001; 94: 153-6.
19. Ballo MS, Sneige N. Can core needle biopsy replace fine needle aspiration cytology in the diagnosis of palpable breast carcinoma: a comparative study of 124 women. *Cancer.* 1996 Aug 15; 78(4): 773-7.
20. Oliver DJ, Frayne JR, Sterrett G. Stereotactic fine needle biopsy of the breast. *Aust N Z J Surg.* 1992; 62: 463-7.
21. Kitchen PRB, Cawson JN. The evolving role of fine needle cytology and core biopsy in the diagnosis of breast cancer. *Aust N Z J Surg.* 1996; 66: 577-9.
22. Frayne J, Sterrett GF, Harvey J, et al. Stereotactic 14 gauge. Core-biopsy of the breast: results from 101 patients. *Aust N Z J Surg.* 1996; 66: 585-91.
23. Florentine BD, Cobb CJ, Frankel K, et al. Core needle biopsy: a useful adjunct to fine needle aspiration in selected patients with palpable breast lesions. *Cancer Cytopathol.* 1997 Feb 25; 81(1): 33-9.
24. Feldman Ps, Covell JL. Breast and lung. In: *Fine needle aspiration cytology and its clinical application.* Chicago-American Society of Clinical Pathologists 1985; p. 27-30.

**Correspondence to: Dr Ali J Awad**

**E-mail: alshuhub2004@yahoo.com**

**Received 29<sup>th</sup> Jan. 2012; Accepted 14<sup>th</sup> Apr. 2012**