

- vitro injection of IFN- γ . *J Immunol.* 1995; 154: 4261–4266.
- 23-** Hill JA, Polgar K and Andreson DJ. T-helper type-1 immunity to trophoblast in women with recurrent spontaneous abortion. *J Am Med Assoc.* 1995; 273: 1933-1936.
- 24-** Makhseed M, Raghupathy R, Azizieh F, et al. Mitogen-induced cytokine responses of maternal lymphocytes suggest a differential Th-type bias in normal pregnancy and pregnancy failure. *Am J Reprod Immunol.* 1999; 42: 273–281.
- 25-** Raghupathy R, Makhseed M. and Azizieh F. Maternal Th1 and Th2 type reactivity to placental antigens in normal and unexplained recurrent Spontaneous abortions. *Cell Immunol.* 1999; 196:122-30.
- 26-** Chaouat G, Menu E, de Smedt D, et al. The emerging role of IL-10 in pregnancy. *Am J Reprod Immunol.* 1996; 35: 325–328.
- 27-** Qstensen M, Sicher P, Förger F and Villiger PM. Activation markers of peripheral blood mononuclear cells in late pregnancy and after delivery:a pilot study. *Ann Rheumat Dis.* 2005; 64:318-20.
- 28-** Marzi M, Vigano A, Tabattoni D, Villa ML, Salvaggio A, et al. Characterization of type 1 and type 2 cytokine profile in physiologic and pathologic human pregnancy. *Clin Exp Immunol.* 1996; 106: 127-133.
- 29-** Kwak-Kim JYH, Chung-Bang HS, Ng SC, Ntrivalas E I, et al. Increased T helper 1 cytokine responses by circulating T cells are present in women with recurrent pregnancy losses and in infertile women with multiple implantation failures after IVF. *Hum Reprod.* 2003; 18:4: 767-773.
- 30-** Saito S, Miyazaki S and Sasaki Y. Th1/Th2 Balance of the implan-tation site in humans. *Immunology of Pregnancy.* 2nd eds. Edited by Mor G. Eureka. Com. 2004; pp. (1-12).
- 31-** Hanna N, Hanna I, Hleb M, Wagner E, et al. Gestational age-dependent expression of IL-10 and its receptor in human placental tissues and isolated cytotrophoblasts. *J Immunol.* 2000; 164(11): 5721 - 5728.
- 32-** Dawood F, Siobhan Q and Fraquharson R. Recurrent Miscarriage: an overview. *Reviews in Gynecological Practice.* 2003; 3: 46-50.
- 33-** Fasouliotis SJ, Spandorfer SD, Witkin SS, Schattman G, et al. Maternal serum levels of interferon- {gamma} and interleukin-2 soluble receptor-{alpha} predict the outcome of early IVF Pregnancies. *Hum Reprod.* 2004; 19(6): 1357 - 1363.
- 34-** Laham N, Brennecke SP and Rice GE Interleukin-8 Release from human gestational tissue explants: effects of gestation,labor, and chorioamnionitis1. *Biol Rreprod.* 1999; 61: 823-827.
- 35-** Laham N, Rice GE, Bishop GJ, Ransome C and Brennecke SP. Interleukin-8 concentrations in amniotic fluid and peripheral venous plasma during human pregnancy and parturition. *Acta Endocrinol* (Copenh). 1993; 129:220-224.
- 36-** Al-Obaidi AB. IFN- γ versus IL-10 In situ expression and immunohistochemistry analysis of VCAM-1, PECAM-1 and HCMV in recurrent spontaneous abortion. M. Sc. Thesis, Coll. Med., Univ. AL-Nahrain. 2005.