

immunoreceptor efficiently lyse target cells in a MHC antigen-and Fas-independent fashion. *J Immunol* 2001; 167: 1090-1096.

16. Iguchi C, Nio Y, Takeda H, et al. Plant polysaccharide PSK: cytostatic effects on growth and invasion; modulating effect on the expression of HLA and adhesion molecules on human gastric and colonic tumor cell surface. *Anticancer Res* 2001; 21: 1007-1013.

17. Tanaka H, Tsunoda T, Nukaya I, et al. Mapping the HLA-A24- restricted T-cell epitope peptide from a tumor-associated antigen HER2/neu: possible immunotherapy for colorectal carcinomas. *B J Cancer* 2001; 84:94-99.

18. Nabeta Y, Sahara H, Suzuki K, et al. Induction of cytotoxic T lymphocytes from peripheral blood of human histocompatibility antigen (HLA)-A31 (+) gastric cancer patients by in vitro stimulation with antigenic peptide of signet ring cell carcinoma. *Jpn J Cancer Res* 2000; 91: 616-621.

19. Schirle M, Keilholz W, Weber B, et al. Identification of tumor-associated MHC class I ligands by a novel T cell-independent approach. *Eur J Immunol* 2000; 30 2216-2225.

20. Novellino PS, Trejo YG, Beviacqua M, et al. Regulation of HLA-DR antigen in monocytes from colorectal cancer patients by in vitro treatment with human recombinant interferon-gamma. *J Invest Allergol Clin Immunol* 2000 10: 90-93.

21. Novellino PS, Trejo YG, Beviacqua M, et al. Cisplatin containing chemotherapy influences HLA-DR expression on monocytes from cancer patients. *J Exp Clin Cancer Res* 1999; 18:481-484.

22. Liu Y, Ng K, O.Lillehei K. Cell mediated immunotherapy: a new approach to the treatment of malignant Glioma. *Cancer Control* 2003; 10(2): 138-147.

23. AH W, WA H, WC L. identification of HLA-A*0201 glioblastoma multiforme cell lines for immunotherapy by PCR-SSP and DNA sequencing. *J Immunooncol.* 2004; 66 (1-2): 1-8

24. Fenstermaker, RA, Ciesielski MJ. Immunotherapeutic strategies for malignant Glioma. *Cancer control.* 2004; 11 (3): 181-191.