

9. Korotzer B, Ong S, Hansen JE. Ethnic differences in pulmonary function in healthy nonsmoking Asian-Americans and European-Americans. *Am J of Respir and Crit Care Med* 2000;161:1101-08.
10. Hankinson JL, Odencrantz JR, Fedan KB. Spirometric reference values from a sample of the general U.S. population. *Am J of Respir and Crit Care Med* 1999;159:179- 87.
11. Quanjer PH, Tammeling GJ, Pederson OF, Peslin R, Yernault JC. Lung volumes and force respiratory flows. Report working party standardization of lung function testing. *Eur Respir J* 1993; 6:5 -30.
12. Knudson RJ, Lebowitz MD, Holberg CJ, Burrows B. Changes in the normal maximal expiratory flow volume curve with growth and aging. *Am Rev Respir Dis* 1983; 127: 725-34.
13. Roca J, Sanchis J, Agusti-Vidal A et al. Spirometric reference values for Mediterranean population. *Bull EUR Physiopathol Respir* 1986; 22: 217-24.
14. Leung, SS, Lau, JT, Xu, YY, et al Secular changes in standing height, sitting height and sexual maturation of Chinese: the Hong Kong Growth Study, 1993. *Ann Hum Biol* 1996; 23:297-306 [Medline]
15. Townsend, MC Spirometric forced expiratory volumes measured in the standing versus the sitting posture. *Am Rev Respir Dis* 1984; 130,123-124 [Medline]
16. Udupihille, M Spirometric and flow standards for healthy adult non-smoking Sri Lankans belonging to the Sinhalese ethnic group. *Ann Hum Bio* 1995; 22,321-336 [Medline]
17. Giri, BR, Sharma, BK, Jindal, SK Normal spirometry in healthy natives of Bhutan. *J Assoc Phys India* 1996; 44,320-322
18. Zheng, JP, Zhong, NS Normative values of pulmonary function testing in Chinese adults. *Chinese Med J* 2002; 2002, 50-54
19. Yang, TS, Peat, J, Keena, V, et al A review of the racial differences in the lung function of normal Caucasian, Chinese and Indian subjects. *Eur Respir J* 1991;4,872-880 [Abstract]
20. Anyanwu CH, Umeh BU. Ventilatory pulmonary function study in healthy young Nigerian adults. *Afr J Med Med Sci* 1989; 18:257-62.
21. Abdullah AK, Abedin MZ, Nouh MS, Al-Nozha M. Ventilatory function in normal Saudi Arabian adults. Observations and comparison with some Western and Eastern reference values. *Trop Geogr Med* 1986; 38:58-62.
22. Verma SS, Kishore N, Raman CV, Lakhera SC, Dass SK. Prediction of some ventilatory 'norms' in healthy Indian males 21-69 years age. *Indian J Physiol Pharmacol* 1983; 27:45-9.
23. Lam KK, Pang SC, Allan WG, Hill LE, Snell NJ, Nunn AJ, et al. A survey of ventilatory capacity in Chinese subjects in Hong Kong. *Ann Hum Biol* 1982; 9:459-72.
24. Fridriksson HV, Malmberg P, Hedenstrom H, Hillerdal G. Reference values for respiratory function tests in males: prediction formulas with tobacco smoking parameters. *Clin Physiol* 1981; 1:349-64.
25. Becklake MR. Concepts of normality applied to the measurement of lung function. *Am J Med* 1986; 80: 1158-64.
26. Schwartz JD, Katz SA, Fegley RW, Tockman MS. Sex and race differences in the development of lung function. *Am Rev Respir Dis* 1988; 138:1415-21.
27. \_uškin E, Smolej-Narančič N, Schachter EN, Mustajbegović J. Spirometric reference values for nonsmoking boys 9-16 years of age. *Acta Med Auxol* 1996; 28: 159-67.
28. Stevens WH, van Hartevelt JH, The PE, Smink HA, Quanjer PH. Validity of ECSC prediction equations for spirometric indices in Dutch conscripts. *Eur Respir J* 1994; 7:29-34.