

wavelength. ⁽¹¹⁾⁽¹⁷⁾⁽¹⁸⁾. The Q-switched Nd: YAG presented better initial as well as long-term results ^{(13) (14)}. Other study done by Jone A, and Grevelink JM, showed that the benefit of Q-switched Nd: YAG laser, it is safe used in darker-skinned patient in whom melanin absorption is concerned ^{(10) (12)}.

In the present study the treatment with Q-switched Nd: YAG laser offers bloodless, low risk, no permanent complication no scarring and no disfigurement in skin was seen, The complication in this study is mainly transient textural change (no scarring) often resolved within one month, and pinpoint bleeding resolved within 5-7 days. There is no change in pigmentation of skin (no hypopigmentation or hyperpigmentation) this probably due to good preparation to patients; sun block is used by patients before and after laser treatment. So it is considered standard treatment for patient seeking tattoo removal.

References

1. Adatto Ma. : In Living Skin – Decorative Tattooing and Dermatology, Editions Roche Basel, (1993).
2. Anderson RR, Geronemus R, Kilmer SL, et al.: Cosmetic tattoo ink darkening. A complication of Q-switched and pulsed- laser treatment. Arch Dermatol. (1993). Aug; 129(8): 1010-4.
3. Apfelberg DB, Maser MR, and Lash H: Comparison of argon and carbon dioxide laser treatment of decorative tattoos: a preliminary report. Ann Plast Surg. (1985): Jan; 14(1): 6-15.
4. Arellano CR, Leopold DA, Shafiroff BB: Tattoo removal comparative study of six methods in the pig. Plast Reconstr Surg (1982). 70(6): 609–703
5. Boo-Chai K: The decorative tattoo: its removal by superficial dermabrasion. Plast Reconstr Surg(1963). 32: 559–563
6. Brady SC, Blokmanis A, and Jewett L: Tattoo removal with the carbon dioxide laser. Ann Plast Surg. (1979): Jun; 2(6): 482-90.
7. Colver GB, Cherry GW, Dawber RP, and Ryan TJ: Tattoo removal using infra-red coagulation. Br J Dermatol (1985). 112(4):481–485
8. Goldman L, Wilson RG, Hornby P, Meyer RG: Radiation from a Q-switched ruby laser. Effect of repeated impacts of power output of 10 megawatt on a tattoo of man. J Invest Dermatol (1965). 44: 69–71
9. Goldstein AP: VII Histologic reactions in tattoos. J Dermatol Surg Oncol(1979). Nov; 5(11): 896-900.
10. Grevelink JM, Duke D, van Leeuwen RL: Laser treatment of tattoos in darkly pigmented patients: efficacy and side effects. J Am Acad Dermatol. (1996): Apr; 34(4): 653-6.
11. Haedersdal M, Bech-Thomsen N, Wulf HC, Hodersdal M: Skin reflectance-guided laser selections for treatment of decorative tattoos. Arch Dermatol (1996): Apr; 132(4): 403-7.
12. Jones A, Roddey P, Orengo I, Rosen T: The Q-switched ND: YAG laser effectively treats tattoos in darkly pigmented skin. Dermatol Surg. Dec; (1996): 22(12): 999-1001.
13. Kaufman, R, Boehncke, WH, Konig, K, Hibst, R: Comparative study of Q-switched Nd: YAG and alexandrite laser treatment of tattoos. Lasers Surg. Med.; (1993). S5: 54.
14. Kaufman R, Hibst R: Pulsed Erbium:YAG laser ablation in cutaneous surgery. Lasers Surg Med; (1996): 19(3): 324-30.
15. Kilmer SL, Lee MS, and Anderson RR: Treatment of multi-colored tattoos with the Q-switched Nd: YAG laser (532 nm): a dose response study with comparison to the Q-switched ruby laser. Lasers Surg Med Suppl. (1993). 5: 54.
16. Kilmer SL, Anderson RR: Clinical use of the Q-switched ruby and the Q-switched Nd: YAG (1064 nm and 532 nm) lasers for treatment of tattoos. J Dermatol Surg Oncol. (1993). 19(4): 330-338.
17. Kilmer SL, Lee MS, Grevelink JM, Flotte TJ, and Anderson RR: The Q-switched Nd: YAG laser effectively treats tattoos. A controlled, dose-response study. Arch Dermatol. (1993). 129:971–978.
18. Kuperman-Beadle M, Levine VJ, Ashinoff R: Laser removal of tattoos. Am J Clin Dermatol. (2001). 2: 21–25.
19. Maurice A. Adatto: Laser: Tattoo Removal: Benefits and Caveat, Med. Laser Appl., (2004). 19: 175–185.
20. Miller TH, Hay-Roe V: Conventional surgery in techniques of removal tattoos. Plast Reconstr Surg 20: 910–914.
21. Piggot TA, Norris RW (1988). : The treatment of tattoos with trichloroacetic acid: