

- men: a hospital-based cross sectional survey, 2004-2006. *Urology*, 2008; 72: 561.
5. Werny DM, Thompson T, Saraiya M, and Freedman D. Obesity is negatively associated with prostate-specific antigen in U.S. men, 2001-2004. *Cancer Epidemiol Biomarkers Prev*, 2007; 16: 70-76.
6. Fowke JH, Signorello LB, Chang SS, Matthews CE, Buchowski MS, Cookson MS, et al. Effects of obesity and height on prostate-specific antigen (PSA) and percentage of free PSA levels among African-American and Caucasian men. *Cancer*, 2006; 107(10): 2361-2367.
7. Randle A, Richards C, Neugut AI. Body composition of abdominal fat distribution and prostate specific antigen test results. *Cancer Epidemiol*, 2009; 18(1): 331-336.
8. Ando R, Nagaya T, Hashimoto Y, Suzuki S, Itoh Y, Umemoto Y, et al. Inverse relationship between obesity and serum prostate-specific antigen level in healthy Japanese men: a hospital-based cross-sectional survey, 2004-2006. *Urology*, 2008; 72(3): 561-565.
9. Chia SE, Lau WK, Chin CM, Tan J, Ho SH, Lee J, et al. Effect of ageing and body mass index on prostate-specific antigen levels among Chinese men in Singapore from a community-based study. *BJU Int*, 2009; 103(11): 1487-1491.
10. Schröder FH, Roobol MJ. Defining the optimal prostate-specific antigen threshold for the diagnosis of prostate cancer. *Curr Opin Urol*, 2009; 19 (3): 227-231.
11. Catalona WJ, Ramos CG, Carvalhal GF, and Yan Y. Lowering PSA cutoffs to enhance detection of curable prostate cancer. *Urology*, 2000; 55: 791-795.
12. Parka JH, Chob BL, Kwon HT, Lee CM, Han HJ. Effect of body mass index and waist circumference on prostate specific antigen and prostate volume in a generally healthy Korean population. *J Urol*; 2010; 182(1): 106-111.
13. Janssen I, Katzmarzyk P, and Ross R. Waist circumference and not body mass index explains obesity-related health risk. *Am J Clin Nutr*, 2004; 79: 379-384.
14. Joseph C, Presti JR, Christopher J, Kane MD, Shinohara K, and Carroll PR. Neoplasm of the prostate gland. *Smith General Urology*, Chapter 22, pp. 360-361.
15. Davies PS, Lanham DA, Stead MA, Tsang K. Body mass index and the prediction of percentage body fat in Australian and Chinese women. *Ann Hum Biol*, 2001; 28: 467-470.
16. Gallagher D, Visser M, Sepulveda D, Pierson RN, Harris T, and Heymsfield SP. How useful is BMI for comparison of body fatness across age, sex, and ethnic groups. *Am J Epidemiol*, 1996; 143: 228-239.
17. Beebe-Dimmer JL, Faerber GJ, Morgenstern H, Werny D, Wojno K, Halstead-Nussloch B, et al. Body composition and serum prostate-specific antigen: review and findings from Flint Men's Health Study. *Urology*, 2008; 71(4): 554-560.
18. Svartberg J, von Muhlen D, Sundsfjord J, and Jorde R. Waist circumference and testosterone levels in community dwelling men. The Tromso study. *Eur J Epidemiol*, 2004; 19: 657-663.
19. Baillargeon J, Pollock BH, and Kristal AR. The association of body mass index and prostate-specific antigen in a population-based study. *Cancer*, 2005; 103: 1092-1095.
20. Porter MP, Stanford JL. Obesity and the risk of prostate cancer. *Prostate*, 2005; 62: 316-321.
21. Bañez LL, Hamilton RJ, Partin AW, Vollmer RT, Sun L, and Rodriguez C. Obesity-related plasma hemodilution and PSA concentration among men with prostate cancer. *Eur Urol*, 2008; 54(1): 232-233.
22. Ohwaki K, Endo F, Muraishi O, Hiramatsu S, and Yano E. Relationship between Prostate-specific Antigen and Hematocrit: Does Hemodilution Lead to Lower PSA Concentrations in Men with a Higher Body Mass Index? *Urology*, 2010 Mar; 75(3): 648-52.
23. Lee SH, Kim JC, Lee JY, Kim JH, Oh CY, Lee SW et al. Effect of obesity on lower urinary tract symptoms in Korean benign prostate hyperplasia patients. *Asian J Androl*, 2009; 11: 663-668.
24. Price MM, Hamilton RJ, Robertson CN, Butts MC, and Freedland SJ. Body mass index, prostate-specific antigen, and digital rectal examination findings among participants in a prostate cancer screening clinic. *Urology*, 2008; 71(5): 787-791.
25. Potosky AL, Feuer EJ and Levin DL. Impact of screening on incidence and mortality of prostate cancer in the united states. *Epidemiol Rev*, 2001; 23: 181.
26. Partin AW, Piantadosi S, and Subong EN. Clearance rate of serum-free and total PSA following radical retropubic prostatectomy. *Prostate*, 1996; 7 Suppl: 35-39.
27. Catalona WJ, Hudson MA, and Scardino PT. Selection of optimal prostate specific antigen cutoffs for early detection of prostate cancer: Receiver operating characteristic curves. *J Urol*, 1994; 152: 2037-2042.

Correspondence to: Dr. Saad D. Farhan,
E-mail: Saadurology@yahoo.com

Received: 18th Feb. 2010, Accepted: 3rd Mar. 2010