

- is a biomarker for EBV-positive Hodgkin's lymphoma. *Clin Cancer Res*, 2006; 12: 460-464.
16. Lo YM, Chan LY, Lo KW, Leung SF, Zhang J, Chan AT, et al. Quantitative analysis of cell free Epstein-Barr virus DNA in plasma of patients with nasopharyngeal carcinoma. *Cancer Res*, 1999; 59: 1188-1191.
 17. Josting A, Franklin J, May M, Koch P, Beykirch MK, Heinz J, et al. New prognostic score based on treatment outcome of patients with relapsed Hodgkin Lymphoma registered in the data base of the German Hodgkins Lymphoma Study Group. *J Clin Oncol*, 2002; 20(1): 221-230.
 18. Schmitz N, and Sureda A. The role of allogeneic stem-cell transplantation in Hodgkin's disease. *Eu J Haematol*, 2005; 75: 146-149.
 19. Fan H, Kim SC, Chima CO, Israel BF, Lawless KM, Eagan PA, Elmore S, et al. Epstein Barr viral load as a marker for lymphoma in AIDS patients. *J Med Virol*, 2004; 75(1): 59-69.
 20. Gallagher A, Armstrong AA, MacKenzie J, Shield L, Khan G, Lake A, et al. Detection of Epstein Barr virus (EBV) genomes in the serum of patients with EBV-associated Hodgkin's disease. *Int J Cancer*, 1999; 84 (4): 942-948.
 21. Drouet E, Brousset P, Fares F, Icart J, Verniol C, Meggetto F, et al. High Epstein-Barr virus serum load and elevated titers of anti ZEBRA antibodies in patients with EBV-harboring tumor cells of Hodgkins disease. *J Med Virol*, 1999; 57(4): 383-389.
 22. Chan KC, Zhang J, Chan AT, Lei KI, Leung SF, Chan LY, et al. Molecular characterization of circulatory EBV DNA in the plasma of nasopharyngeal carcinoma and lymphoma patients. *Cancer Res*, 2003; 63(9): 2028-2032.
 23. Au WY, Pang A, Choy C, Chim CS, Kwong YL. Quantification of circulating Epstein-Barr Virus (EBV) DNA in the diagnosis and monitoring of natural killer cell and EBV positive lymphoma in immunocompetent patients. *Blood*, 2004; 104(1): 243-249.
 24. Lin JC, Wang WY, Chen KY, Wei YH, Liang WM, Jan JS, et al. Quantification of plasma Epstein-Barr virus DNA in patients with advanced nasopharyngeal carcinoma. *N Eng J*, 2004; 350: 2461-2470.
 25. Fan H, and Gullay ML. Epstein-Barr viral load measurement as a marker of EBV related disease. *Mol Diagnostic*, 2001; 6: 279-288.
 26. Chan KC, Zhang J, Chan AT, Lei KI, Leung SF, Chan LY, et al. Molecular characterization of circulatory EBV DNA in the plasma of nasopharyngeal carcinoma and lymphoma patients. *Cancer Res*, 2003; 63(9): 2028-2032.
 27. Jahr S, Hentze H, Englisch S, Hardt D, Fackelmayer FO, Hesch RD, et al. DNA fragments in the blood plasma of cancer patients: quantification and evidence of their origin from apoptotic and necrotic cells. *Cancer Res*, 2001; 61(4): 1659-1665.

Correspondence to: Dr. Elham A. Alaswad

E-mail: elham_alaswad@yahoo.com

Received: 31st May 2009, Accepted: 16th Dec. 2009.